

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -7 & -7 & 1 \\ 5 & 5 & -1 \\ -3 & -3 & 1 \end{pmatrix} & (b) \begin{pmatrix} -3 & 4 & 4 \\ 0 & -3 & 0 \\ 0 & 2 & -1 \end{pmatrix} & (c) \begin{pmatrix} -2 & -6 & 3 \\ 4 & 0 & 5 \\ 2 & 4 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 2 & 2 \\ -1 & -4 & -1 \\ -1 & -1 & -4 \end{pmatrix} & (e) \begin{pmatrix} -1 & 1 & 2 \\ 2 & -2 & 5 \\ -2 & -1 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -7 & -7 & 6 \\ 8 & 8 & -6 \\ -5 & -5 & 6 \end{pmatrix} & (b) \begin{pmatrix} 0 & 0 & 3 \\ -2 & -2 & -3 \\ -2 & 0 & -5 \end{pmatrix} & (c) \begin{pmatrix} -4 & -8 & -7 \\ 1 & -3 & 1 \\ 3 & 7 & 6 \end{pmatrix} \\ (d) \begin{pmatrix} 1 & -3 & 6 \\ -1 & -1 & -2 \\ -2 & 2 & -6 \end{pmatrix} & (e) \begin{pmatrix} 0 & 2 & 4 \\ -1 & 0 & 1 \\ -2 & -2 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -7 & -6 & -9 \\ 4 & 5 & 4 \\ 6 & 4 & 8 \end{pmatrix} \quad (b) \begin{pmatrix} -7 & 5 & -5 \\ -5 & 3 & -5 \\ 5 & -5 & 3 \end{pmatrix} \quad (c) \begin{pmatrix} -3 & -2 & 3 \\ -2 & -3 & 3 \\ -4 & -8 & 7 \end{pmatrix}$$
$$(d) \begin{pmatrix} -4 & 2 & -2 \\ 0 & -2 & 0 \\ 2 & -2 & 0 \end{pmatrix} \quad (e) \begin{pmatrix} -4 & -2 & 6 \\ -4 & -6 & 6 \\ -3 & -3 & 4 \end{pmatrix}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -7 & -6 & -7 \\ -6 & -3 & -6 \\ 2 & 6 & 2 \end{pmatrix} & (b) \begin{pmatrix} 1 & 0 & -4 \\ 2 & -3 & -2 \\ 2 & 0 & -5 \end{pmatrix} & (c) \begin{pmatrix} -1 & 0 & 2 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 1 & 0 \\ -1 & -3 & 0 \\ 1 & 1 & -2 \end{pmatrix} & (e) \begin{pmatrix} -6 & 4 & 4 \\ -2 & 0 & 2 \\ -1 & 0 & 0 \end{pmatrix} & \end{array}$$

- [2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.
- [3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -7 & -6 & 5 \\ 7 & 6 & -5 \\ -4 & -4 & 4 \end{pmatrix} \quad (b) \begin{pmatrix} -5 & 0 & 4 \\ -2 & -1 & 2 \\ -2 & 0 & 1 \end{pmatrix} \quad (c) \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 1 \\ -1 & -1 & 2 \end{pmatrix}$$
$$(d) \begin{pmatrix} -1 & 2 & -2 \\ 0 & -2 & 1 \\ 0 & -1 & 0 \end{pmatrix} \quad (e) \begin{pmatrix} 4 & 0 & -6 \\ 3 & -2 & -3 \\ 8 & -4 & -8 \end{pmatrix}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁶

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -7 & -3 & -5 \\ 7 & 3 & 5 \\ -1 & -1 & 1 \end{pmatrix} \quad (b) \begin{pmatrix} -6 & -2 & 4 \\ 6 & 1 & -6 \\ -2 & -1 & 0 \end{pmatrix} \quad (c) \begin{pmatrix} 3 & -1 & -1 \\ 0 & 2 & -1 \\ 1 & -1 & 3 \end{pmatrix}$$

$$(d) \begin{pmatrix} -3 & 4 & 0 \\ -1 & 1 & 0 \\ -2 & 4 & -1 \end{pmatrix} \quad (e) \begin{pmatrix} 2 & -1 & -3 \\ -1 & 0 & 1 \\ 4 & -2 & -5 \end{pmatrix}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁶Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁷

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -6 & -8 & 5 \\ 7 & 9 & -5 \\ -6 & -6 & 7 \end{pmatrix} & (b) \begin{pmatrix} -4 & 6 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} & (c) \begin{pmatrix} 4 & -1 & -1 \\ -1 & 4 & 1 \\ 4 & -5 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} -3 & 4 & 0 \\ -1 & 1 & 0 \\ -3 & 6 & -1 \end{pmatrix} & (e) \begin{pmatrix} 0 & 1 & 0 \\ -1 & 2 & 0 \\ 2 & -3 & 1 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁷Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁸

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -6 & -6 & 3 \\ 5 & 5 & -3 \\ -4 & -4 & 3 \end{pmatrix} & (b) \begin{pmatrix} 7 & -8 & -4 \\ 0 & -1 & 0 \\ 8 & -8 & -5 \end{pmatrix} & (c) \begin{pmatrix} 2 & 0 & 3 \\ 3 & -1 & 3 \\ 6 & -6 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & -8 & 4 \\ 0 & -5 & 2 \\ 0 & -8 & 3 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & 0 \\ -2 & 1 & -1 \\ 2 & 3 & 3 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁸Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -6 & -5 & 4 \\ 6 & 5 & -4 \\ -3 & -3 & 3 \end{pmatrix} \quad (b) \begin{pmatrix} -2 & 1 & -3 \\ 1 & -2 & 3 \\ 1 & -1 & 2 \end{pmatrix} S \quad (c) \begin{pmatrix} -2 & -6 & 3 \\ 4 & 0 & 5 \\ 2 & 4 & -1 \end{pmatrix}$$
$$(d) \begin{pmatrix} 3 & -4 & 0 \\ 1 & -1 & 0 \\ 2 & -4 & 1 \end{pmatrix} \quad (e) \begin{pmatrix} -4 & -6 & -3 \\ 8 & 8 & 4 \\ 0 & 3 & 2 \end{pmatrix}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹⁰

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -6 & 0 & -2 \\ -3 & 0 & -3 \\ 3 & 0 & -1 \end{pmatrix} & (b) \begin{pmatrix} -1 & -2 & 6 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} & (c) \begin{pmatrix} -4 & -8 & -7 \\ 1 & -3 & 1 \\ 3 & 7 & 6 \end{pmatrix} \\ (d) \begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & -6 \\ 0 & 0 & 1 \end{pmatrix} & (e) \begin{pmatrix} 4 & -1 & -1 \\ 2 & -1 & -2 \\ -3 & 7 & 6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹⁰Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹¹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -6 & 5 & -6 \\ -6 & 5 & -6 \\ 4 & -4 & 2 \end{pmatrix} \quad (b) \begin{pmatrix} -1 & 2 & 0 \\ -1 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (c) \begin{pmatrix} -3 & -2 & 3 \\ -2 & -3 & 3 \\ -4 & -8 & 7 \end{pmatrix}$$

$$(d) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & -6 & 1 \end{pmatrix} \quad (e) \begin{pmatrix} 2 & 2 & 1 \\ 3 & 0 & -3 \\ -4 & 5 & 7 \end{pmatrix}$$

- [2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.
- [3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹¹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹²

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -6 & 8 & -6 \\ -8 & 9 & -8 \\ 3 & -4 & 3 \end{pmatrix} & (b) \begin{pmatrix} -3 & -8 & 8 \\ 0 & 1 & -4 \\ 0 & 0 & -3 \end{pmatrix} & (c) \begin{pmatrix} -1 & 0 & 2 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} 2 & -8 & -4 \\ 0 & -2 & -2 \\ 0 & 8 & 6 \end{pmatrix} & (e) \begin{pmatrix} -1 & 1 & 2 \\ 2 & -2 & 5 \\ -2 & -1 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹²Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹³

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -5 & -9 & 2 \\ 2 & 6 & -2 \\ -5 & -5 & 2 \end{pmatrix} & (b) \begin{pmatrix} 1 & -6 & -3 \\ 0 & -1 & -1 \\ 0 & 2 & 2 \end{pmatrix} & (c) \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 1 \\ -1 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} 2 & 1 & 2 \\ 1 & 2 & -2 \\ -1 & 1 & 5 \end{pmatrix} & (e) \begin{pmatrix} 0 & 2 & 4 \\ -1 & 0 & 1 \\ -2 & -2 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹³Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹⁴

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -5 & -7 & 1 \\ 5 & 7 & -1 \\ -3 & -3 & 3 \end{pmatrix} & (b) \begin{pmatrix} 2 & 2 & 0 \\ 0 & 1 & 0 \\ -2 & -4 & 1 \end{pmatrix} & (c) \begin{pmatrix} 3 & -1 & -1 \\ 0 & 2 & -1 \\ 1 & -1 & 3 \end{pmatrix} \\ (d) \begin{pmatrix} 7 & -4 & -2 \\ 4 & -1 & -2 \\ 0 & 0 & 3 \end{pmatrix} & (e) \begin{pmatrix} -4 & -2 & 6 \\ -4 & -6 & 6 \\ -3 & -3 & 4 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹⁴Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹⁵

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -5 & -6 & 0 \\ 4 & -3 & 4 \\ 6 & 6 & 1 \end{pmatrix} & (b) \begin{pmatrix} -2 & 4 & 0 \\ 0 & 2 & 0 \\ -4 & 4 & 2 \end{pmatrix} & (c) \begin{pmatrix} 4 & -1 & -1 \\ -1 & 4 & 1 \\ 4 & -5 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 2 & 2 \\ -1 & -4 & -1 \\ -1 & -1 & -4 \end{pmatrix} & (e) \begin{pmatrix} -6 & 4 & 4 \\ -2 & 0 & 2 \\ -1 & 0 & 0 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹⁵Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹⁶

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -5 & -3 & 3 \\ 2 & 0 & -2 \\ -6 & -6 & 2 \end{pmatrix} \quad (b) \begin{pmatrix} 2 & 0 & 0 \\ 8 & -2 & -4 \\ -4 & 2 & 4 \end{pmatrix} \quad (c) \begin{pmatrix} 2 & 0 & 3 \\ 3 & -1 & 3 \\ 6 & -6 & 2 \end{pmatrix}$$
$$(d) \begin{pmatrix} 1 & -3 & 6 \\ -1 & -1 & -2 \\ -2 & 2 & -6 \end{pmatrix} \quad (e) \begin{pmatrix} 4 & 0 & -6 \\ 3 & -2 & -3 \\ 8 & -4 & -8 \end{pmatrix}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹⁶Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹⁷

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -5 & 0 & -2 \\ -3 & 1 & -3 \\ 3 & 0 & 0 \end{pmatrix} & (b) \begin{pmatrix} 1 & -1 & 0 \\ 0 & 2 & 0 \\ -1 & -1 & 2 \end{pmatrix} & (c) \begin{pmatrix} -2 & -6 & 3 \\ 4 & 0 & 5 \\ 2 & 4 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} -4 & 2 & -2 \\ 0 & -2 & 0 \\ 2 & -2 & 0 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & -3 \\ -1 & 0 & 1 \\ 4 & -2 & -5 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹⁷Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹⁸

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

(a) $\begin{pmatrix} -5 & 5 & 5 \\ -2 & 4 & 3 \\ -8 & 6 & 7 \end{pmatrix}$

(b) $\begin{pmatrix} 6 & -4 & 1 \\ 3 & -1 & 1 \\ 0 & 0 & 3 \end{pmatrix}$

(c) $\begin{pmatrix} -4 & -8 & -7 \\ 1 & -3 & 1 \\ 3 & 7 & 6 \end{pmatrix}$

(d) $\begin{pmatrix} -1 & 1 & 0 \\ -1 & -3 & 0 \\ 1 & 1 & -2 \end{pmatrix}$

(e) $\begin{pmatrix} 0 & 1 & 0 \\ -1 & 2 & 0 \\ 2 & -3 & 1 \end{pmatrix}$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹⁸Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work¹⁹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -5 & 6 & -4 \\ -6 & 6 & -6 \\ 2 & -3 & 1 \end{pmatrix} & (b) \begin{pmatrix} 3 & -2 & -2 \\ 0 & 3 & 0 \\ 0 & -2 & 1 \end{pmatrix} & (c) \begin{pmatrix} -3 & -2 & 3 \\ -2 & -3 & 3 \\ -4 & -8 & 7 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 2 & -2 \\ 0 & -2 & 1 \\ 0 & -1 & 0 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & 0 \\ -2 & 1 & -1 \\ 2 & 3 & 3 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

¹⁹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²⁰

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & -6 & 3 \\ 5 & 7 & -3 \\ -4 & -4 & 5 \end{pmatrix} & (b) \begin{pmatrix} 5 & 0 & 2 \\ 2 & 3 & 2 \\ -4 & 0 & -1 \end{pmatrix} & (c) \begin{pmatrix} -1 & 0 & 2 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -3 & 4 & 0 \\ -1 & 1 & 0 \\ -2 & 4 & -1 \end{pmatrix} & (e) \begin{pmatrix} -4 & -6 & -3 \\ 8 & 8 & 4 \\ 0 & 3 & 2 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²⁰Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²¹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & -5 & 3 \\ 7 & 8 & -3 \\ 5 & 5 & -2 \end{pmatrix} & (b) \begin{pmatrix} 1 & -1 & 0 \\ 2 & 4 & 0 \\ 4 & 2 & 3 \end{pmatrix} & (c) \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 1 \\ -1 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -3 & 4 & 0 \\ -1 & 1 & 0 \\ -3 & 6 & -1 \end{pmatrix} & (e) \begin{pmatrix} 4 & -1 & -1 \\ 2 & -1 & -2 \\ -3 & 7 & 6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²¹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²²

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & -4 & -2 \\ 7 & 7 & 2 \\ 1 & 1 & 2 \end{pmatrix} & (b) \begin{pmatrix} -3 & 4 & 4 \\ 0 & -3 & 0 \\ 0 & 2 & -1 \end{pmatrix} & (c) \begin{pmatrix} 3 & -1 & -1 \\ 0 & 2 & -1 \\ 1 & -1 & 3 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & -8 & 4 \\ 0 & -5 & 2 \\ 0 & -8 & 3 \end{pmatrix} & (e) \begin{pmatrix} 2 & 2 & 1 \\ 3 & 0 & -3 \\ -4 & 5 & 7 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²²Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²³

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & -4 & 1 \\ -3 & -3 & 3 \\ -6 & -6 & 3 \end{pmatrix} & (b) \begin{pmatrix} 0 & 0 & 3 \\ -2 & -2 & -3 \\ -2 & 0 & -5 \end{pmatrix} & (c) \begin{pmatrix} 4 & -1 & -1 \\ -1 & 4 & 1 \\ 4 & -5 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} 3 & -4 & 0 \\ 1 & -1 & 0 \\ 2 & -4 & 1 \end{pmatrix} & (e) \begin{pmatrix} -1 & 1 & 2 \\ 2 & -2 & 5 \\ -2 & -1 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²³Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²⁴

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & -4 & 5 \\ 3 & 3 & -5 \\ 4 & 4 & -5 \end{pmatrix} & (b) \begin{pmatrix} -7 & 5 & -5 \\ -5 & 3 & -5 \\ 5 & -5 & 3 \end{pmatrix} & (c) \begin{pmatrix} 2 & 0 & 3 \\ 3 & -1 & 3 \\ 6 & -6 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & -6 \\ 0 & 0 & 1 \end{pmatrix} & (e) \begin{pmatrix} 0 & 2 & 4 \\ -1 & 0 & 1 \\ -2 & -2 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²⁴Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²⁵

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & -3 & -1 \\ -2 & 3 & -2 \\ 2 & 6 & -1 \end{pmatrix} & (b) \begin{pmatrix} 1 & 0 & -4 \\ 2 & -3 & -2 \\ 2 & 0 & -5 \end{pmatrix} & (c) \begin{pmatrix} -2 & -6 & 3 \\ 4 & 0 & 5 \\ 2 & 4 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & -6 & 1 \end{pmatrix} & (e) \begin{pmatrix} -4 & -2 & 6 \\ -4 & -6 & 6 \\ -3 & -3 & 4 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²⁵Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²⁶

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & -2 & -5 \\ 2 & 0 & 2 \\ 2 & 2 & 3 \end{pmatrix} & (b) \begin{pmatrix} -5 & 0 & 4 \\ -2 & -1 & 2 \\ -2 & 0 & 1 \end{pmatrix} & (c) \begin{pmatrix} -4 & -8 & -7 \\ 1 & -3 & 1 \\ 3 & 7 & 6 \end{pmatrix} \\ (d) \begin{pmatrix} 2 & -8 & -4 \\ 0 & -2 & -2 \\ 0 & 8 & 6 \end{pmatrix} & (e) \begin{pmatrix} -6 & 4 & 4 \\ -2 & 0 & 2 \\ -1 & 0 & 0 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²⁶Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²⁷

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & -1 & -3 \\ 3 & 0 & 3 \\ -1 & -1 & 0 \end{pmatrix} & (b) \begin{pmatrix} -6 & -2 & 4 \\ 6 & 1 & -6 \\ -2 & -1 & 0 \end{pmatrix} & (c) \begin{pmatrix} -3 & -2 & 3 \\ -2 & -3 & 3 \\ -4 & -8 & 7 \end{pmatrix} \\ (d) \begin{pmatrix} 2 & 1 & 2 \\ 1 & 2 & -2 \\ -1 & 1 & 5 \end{pmatrix} & (e) \begin{pmatrix} 4 & 0 & -6 \\ 3 & -2 & -3 \\ 8 & -4 & -8 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²⁷Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²⁸

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & -1 & 2 \\ -4 & -1 & 2 \\ -5 & 1 & 1 \end{pmatrix} & (b) \begin{pmatrix} -4 & 6 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} & (c) \begin{pmatrix} -1 & 0 & 2 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} 7 & -4 & -2 \\ 4 & -1 & -2 \\ 0 & 0 & 3 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & -3 \\ -1 & 0 & 1 \\ 4 & -2 & -5 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²⁸Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work²⁹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & 0 & -6 \\ 4 & 0 & 6 \\ 1 & 0 & 1 \end{pmatrix} & (b) \begin{pmatrix} 7 & -8 & -4 \\ 0 & -1 & 0 \\ 8 & -8 & -5 \end{pmatrix} & (c) \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 1 \\ -1 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 2 & 2 \\ -1 & -4 & -1 \\ -1 & -1 & -4 \end{pmatrix} & (e) \begin{pmatrix} 0 & 1 & 0 \\ -1 & 2 & 0 \\ 2 & -3 & 1 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

²⁹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³⁰

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & 2 & 2 \\ -4 & 2 & 2 \\ 4 & -4 & -4 \end{pmatrix} & (b) \begin{pmatrix} -2 & 1 & -3 \\ 1 & -2 & 3 \\ 1 & -1 & 2 \end{pmatrix} & (c) \begin{pmatrix} 3 & -1 & -1 \\ 0 & 2 & -1 \\ 1 & -1 & 3 \end{pmatrix} \\ (d) \begin{pmatrix} 1 & -3 & 6 \\ -1 & -1 & -2 \\ -2 & 2 & -6 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & 0 \\ -2 & 1 & -1 \\ 2 & 3 & 3 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³⁰Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³¹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & 4 & 4 \\ -7 & 7 & 4 \\ 7 & -7 & -4 \end{pmatrix} & (b) \begin{pmatrix} -1 & -2 & 6 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} & (c) \begin{pmatrix} 4 & -1 & -1 \\ -1 & 4 & 1 \\ 4 & -5 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} -4 & 2 & -2 \\ 0 & -2 & 0 \\ 2 & -2 & 0 \end{pmatrix} & (e) \begin{pmatrix} -4 & -6 & -3 \\ 8 & 8 & 4 \\ 0 & 3 & 2 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³¹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³²

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & 4 & 4 \\ -5 & 5 & 4 \\ 5 & -5 & -4 \end{pmatrix} & (b) \begin{pmatrix} -1 & 2 & 0 \\ -1 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix} & (c) \begin{pmatrix} 2 & 0 & 3 \\ 3 & -1 & 3 \\ 6 & -6 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 1 & 0 \\ -1 & -3 & 0 \\ 1 & 1 & -2 \end{pmatrix} & (e) \begin{pmatrix} 4 & -1 & -1 \\ 2 & -1 & -2 \\ -3 & 7 & 6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³²Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³³

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & 6 & 6 \\ -2 & 4 & 5 \\ -2 & 2 & 1 \end{pmatrix} & (b) \begin{pmatrix} -3 & -8 & 8 \\ 0 & 1 & -4 \\ 0 & 0 & -3 \end{pmatrix} & (c) \begin{pmatrix} -2 & -6 & 3 \\ 4 & 0 & 5 \\ 2 & 4 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 2 & -2 \\ 0 & -2 & 1 \\ 0 & -1 & 0 \end{pmatrix} & (e) \begin{pmatrix} 2 & 2 & 1 \\ 3 & 0 & -3 \\ -4 & 5 & 7 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³³Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³⁴

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -4 & 6 & 6 \\ 2 & 0 & 3 \\ -6 & 6 & 3 \end{pmatrix} & (b) \begin{pmatrix} 1 & -6 & -3 \\ 0 & -1 & -1 \\ 0 & 2 & 2 \end{pmatrix} & (c) \begin{pmatrix} -4 & -8 & -7 \\ 1 & -3 & 1 \\ 3 & 7 & 6 \end{pmatrix} \\ (d) \begin{pmatrix} -3 & 4 & 0 \\ -1 & 1 & 0 \\ -2 & 4 & -1 \end{pmatrix} & (e) \begin{pmatrix} -1 & 1 & 2 \\ 2 & -2 & 5 \\ -2 & -1 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³⁴Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³⁵

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -3 & -7 & 1 \\ 1 & 5 & -1 \\ -5 & -5 & 3 \end{pmatrix} & (b) \begin{pmatrix} 2 & 2 & 0 \\ 0 & 1 & 0 \\ -2 & -4 & 1 \end{pmatrix} & (c) \begin{pmatrix} -3 & -2 & 3 \\ -2 & -3 & 3 \\ -4 & -8 & 7 \end{pmatrix} \\ (d) \begin{pmatrix} -3 & 4 & 0 \\ -1 & 1 & 0 \\ -3 & 6 & -1 \end{pmatrix} & (e) \begin{pmatrix} 0 & 2 & 4 \\ -1 & 0 & 1 \\ -2 & -2 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³⁵Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³⁶

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -3 & -6 & -4 \\ -5 & -2 & -5 \\ 6 & 6 & 7 \end{pmatrix} & (b) \begin{pmatrix} -2 & 4 & 0 \\ 0 & 2 & 0 \\ -4 & 4 & 2 \end{pmatrix} & (c) \begin{pmatrix} -1 & 0 & 2 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & -8 & 4 \\ 0 & -5 & 2 \\ 0 & -8 & 3 \end{pmatrix} & (e) \begin{pmatrix} -4 & -2 & 6 \\ -4 & -6 & 6 \\ -3 & -3 & 4 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³⁶Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³⁷

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -3 & -4 & -6 \\ 5 & 6 & 6 \\ 5 & 5 & 7 \end{pmatrix} \quad (b) \begin{pmatrix} 2 & 0 & 0 \\ 8 & -2 & -4 \\ -4 & 2 & 4 \end{pmatrix} \quad (c) \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 1 \\ -1 & -1 & 2 \end{pmatrix}$$
$$(d) \begin{pmatrix} 3 & -4 & 0 \\ 1 & -1 & 0 \\ 2 & -4 & 1 \end{pmatrix} \quad (e) \begin{pmatrix} -6 & 4 & 4 \\ -2 & 0 & 2 \\ -1 & 0 & 0 \end{pmatrix}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³⁷Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³⁸

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -3 & -4 & -5 \\ -1 & 0 & -1 \\ 4 & 4 & 6 \end{pmatrix} & (b) \begin{pmatrix} 1 & -1 & 0 \\ 0 & 2 & 0 \\ -1 & -1 & 2 \end{pmatrix} & (c) \begin{pmatrix} 3 & -1 & -1 \\ 0 & 2 & -1 \\ 1 & -1 & 3 \end{pmatrix} \\ (d) \begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & -6 \\ 0 & 0 & 1 \end{pmatrix} & (e) \begin{pmatrix} 4 & 0 & -6 \\ 3 & -2 & -3 \\ 8 & -4 & -8 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³⁸Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work³⁹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -3 & -3 & -5 \\ 4 & 4 & 5 \\ 4 & 4 & 5 \end{pmatrix} & (b) \begin{pmatrix} 6 & -4 & 1 \\ 3 & -1 & 1 \\ 0 & 0 & 3 \end{pmatrix} & (c) \begin{pmatrix} 4 & -1 & -1 \\ -1 & 4 & 1 \\ 4 & -5 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & -6 & 1 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & -3 \\ -1 & 0 & 1 \\ 4 & -2 & -5 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

³⁹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴⁰

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -3 & -3 & 0 \\ 4 & 2 & 2 \\ 3 & -3 & 6 \end{pmatrix} & (b) \begin{pmatrix} 3 & -2 & -2 \\ 0 & 3 & 0 \\ 0 & -2 & 1 \end{pmatrix} & (c) \begin{pmatrix} 2 & 0 & 3 \\ 3 & -1 & 3 \\ 6 & -6 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} 2 & -8 & -4 \\ 0 & -2 & -2 \\ 0 & 8 & 6 \end{pmatrix} & (e) \begin{pmatrix} 0 & 1 & 0 \\ -1 & 2 & 0 \\ 2 & -3 & 1 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴⁰Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴¹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -3 & 0 & -8 \\ 1 & -2 & 8 \\ 4 & 0 & 9 \end{pmatrix} & (b) \begin{pmatrix} 5 & 0 & 2 \\ 2 & 3 & 2 \\ -4 & 0 & -1 \end{pmatrix} & (c) \begin{pmatrix} -2 & -6 & 3 \\ 4 & 0 & 5 \\ 2 & 4 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} 2 & 1 & 2 \\ 1 & 2 & -2 \\ -1 & 1 & 5 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & 0 \\ -2 & 1 & -1 \\ 2 & 3 & 3 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴¹Date: Dec 19, 2009; Dead Line : 27 Dec 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴²

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -3 & 1 & 1 \\ 5 & -3 & -3 \\ -7 & 3 & 3 \end{pmatrix} & (b) \begin{pmatrix} 1 & -1 & 0 \\ 2 & 4 & 0 \\ 4 & 2 & 3 \end{pmatrix} & (c) \begin{pmatrix} -4 & -8 & -7 \\ 1 & -3 & 1 \\ 3 & 7 & 6 \end{pmatrix} \\ (d) \begin{pmatrix} 7 & -4 & -2 \\ 4 & -1 & -2 \\ 0 & 0 & 3 \end{pmatrix} & (e) \begin{pmatrix} -4 & -6 & -3 \\ 8 & 8 & 4 \\ 0 & 3 & 2 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴²Date: Dec 19, 2009; Dead Line : 27 Dec 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴³

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -3 & 5 & 5 \\ 6 & -4 & 1 \\ -6 & 6 & 1 \end{pmatrix} & (b) \begin{pmatrix} -3 & 4 & 4 \\ 0 & -3 & 0 \\ 0 & 2 & -1 \end{pmatrix} & (c) \begin{pmatrix} -3 & -2 & 3 \\ -2 & -3 & 3 \\ -4 & -8 & 7 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 2 & 2 \\ -1 & -4 & -1 \\ -1 & -1 & -4 \end{pmatrix} & (e) \begin{pmatrix} 4 & -1 & -1 \\ 2 & -1 & -2 \\ -3 & 7 & 6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴³Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴⁴

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -3 & 6 & -2 \\ 4 & 2 & 4 \\ 7 & -6 & 6 \end{pmatrix} & (b) \begin{pmatrix} 0 & 0 & 3 \\ -2 & -2 & -3 \\ -2 & 0 & -5 \end{pmatrix} & (c) \begin{pmatrix} -1 & 0 & 2 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} 1 & -3 & 6 \\ -1 & -1 & -2 \\ -2 & 2 & -6 \end{pmatrix} & (e) \begin{pmatrix} 2 & 2 & 1 \\ 3 & 0 & -3 \\ -4 & 5 & 7 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴⁴Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴⁵

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -2 & -5 & 1 \\ 1 & 4 & -1 \\ -3 & -3 & 2 \end{pmatrix} & (b) \begin{pmatrix} -7 & 5 & -5 \\ -5 & 3 & -5 \\ 5 & -5 & 3 \end{pmatrix} & (c) \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 1 \\ -1 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -4 & 2 & -2 \\ 0 & -2 & 0 \\ 2 & -2 & 0 \end{pmatrix} & (e) \begin{pmatrix} -1 & 1 & 2 \\ 2 & -2 & 5 \\ -2 & -1 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴⁵Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴⁶

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -2 & -3 & 1 \\ 5 & 6 & -1 \\ 3 & 3 & 0 \end{pmatrix} & (b) \begin{pmatrix} 1 & 0 & -4 \\ 2 & -3 & -2 \\ 2 & 0 & -5 \end{pmatrix} & (c) \begin{pmatrix} 3 & -1 & -1 \\ 0 & 2 & -1 \\ 1 & -1 & 3 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 1 & 0 \\ -1 & -3 & 0 \\ 1 & 1 & -2 \end{pmatrix} & (e) \begin{pmatrix} 0 & 2 & 4 \\ -1 & 0 & 1 \\ -2 & -2 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴⁶Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴⁷

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -2 & -1 & -5 \\ 4 & 3 & 5 \\ -2 & -2 & 4 \end{pmatrix}$$

$$(b) \begin{pmatrix} -5 & 0 & 4 \\ -2 & -1 & 2 \\ -2 & 0 & 1 \end{pmatrix}$$

$$(c) \begin{pmatrix} 4 & -1 & -1 \\ -1 & 4 & 1 \\ 4 & -5 & -1 \end{pmatrix}$$

$$(d) \begin{pmatrix} -1 & 2 & -2 \\ 0 & -2 & 1 \\ 0 & -1 & 0 \end{pmatrix}$$

$$(e) \begin{pmatrix} -4 & -2 & 6 \\ -4 & -6 & 6 \\ -3 & -3 & 4 \end{pmatrix}$$

- [2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.
- [3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴⁷Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴⁸

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -2 & 0 & -6 \\ 0 & -2 & 6 \\ 3 & 0 & 7 \end{pmatrix} \quad (b) \begin{pmatrix} -6 & -2 & 4 \\ 6 & 1 & -6 \\ -2 & -1 & 0 \end{pmatrix} \quad (c) \begin{pmatrix} 2 & 0 & 3 \\ 3 & -1 & 3 \\ 6 & -6 & 2 \end{pmatrix}$$

$$(d) \begin{pmatrix} -3 & 4 & 0 \\ -1 & 1 & 0 \\ -2 & 4 & -1 \end{pmatrix} \quad (e) \begin{pmatrix} -6 & 4 & 4 \\ -2 & 0 & 2 \\ -1 & 0 & 0 \end{pmatrix}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴⁸Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁴⁹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -2 & 0 & -6 \\ 3 & 1 & 6 \\ 3 & 3 & 4 \end{pmatrix} & (b) \begin{pmatrix} -4 & 6 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} & (c) \begin{pmatrix} -2 & -6 & 3 \\ 4 & 0 & 5 \\ 2 & 4 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} -3 & 4 & 0 \\ -1 & 1 & 0 \\ -3 & 6 & -1 \end{pmatrix} & (e) \begin{pmatrix} 4 & 0 & -6 \\ 3 & -2 & -3 \\ 8 & -4 & -8 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁴⁹Date: Dec 19, 2009; Dead Line : 27 Dec 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵⁰

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -2 & 0 & -4 \\ -1 & -3 & 4 \\ 2 & 0 & 4 \end{pmatrix} & (b) \begin{pmatrix} 7 & -8 & -4 \\ 0 & -1 & 0 \\ 8 & -8 & -5 \end{pmatrix} & (c) \begin{pmatrix} -4 & -8 & -7 \\ 1 & -3 & 1 \\ 3 & 7 & 6 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & -8 & 4 \\ 0 & -5 & 2 \\ 0 & -8 & 3 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & -3 \\ -1 & 0 & 1 \\ 4 & -2 & -5 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵⁰Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵¹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -2 & 0 & 3 \\ -6 & 4 & 3 \\ -2 & 0 & 3 \end{pmatrix}$$

$$(b) \begin{pmatrix} -2 & 1 & -3 \\ 1 & -2 & 3 \\ 1 & -1 & 2 \end{pmatrix}$$

$$(c) \begin{pmatrix} -3 & -2 & 3 \\ -2 & -3 & 3 \\ -4 & -8 & 7 \end{pmatrix}$$

$$(d) \begin{pmatrix} 3 & -4 & 0 \\ 1 & -1 & 0 \\ 2 & -4 & 1 \end{pmatrix}$$

$$(e) \begin{pmatrix} 0 & 1 & 0 \\ -1 & 2 & 0 \\ 2 & -3 & 1 \end{pmatrix}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵¹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵²

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$(a) \begin{pmatrix} -2 & 0 & 3 \\ 0 & -2 & 6 \\ -2 & 0 & 3 \end{pmatrix} \quad (b) \begin{pmatrix} -1 & -2 & 6 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} \quad (c) \begin{pmatrix} -1 & 0 & 2 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix}$$

$$(d) \begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & -6 \\ 0 & 0 & 1 \end{pmatrix} \quad (e) \begin{pmatrix} 2 & -1 & 0 \\ -2 & 1 & -1 \\ 2 & 3 & 3 \end{pmatrix}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵²Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵³

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -2 & 0 & 3 \\ 0 & -2 & 6 \\ -2 & 0 & 3 \end{pmatrix} & (b) \begin{pmatrix} -1 & -2 & 6 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} & (c) \begin{pmatrix} -1 & 0 & 2 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & -6 \\ 0 & 0 & 1 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & 0 \\ -2 & 1 & -1 \\ 2 & 3 & 3 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵³Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵⁴

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -2 & 2 & -6 \\ 2 & -2 & 6 \\ 2 & 2 & 2 \end{pmatrix} & (b) \begin{pmatrix} -3 & -8 & 8 \\ 0 & 1 & -4 \\ 0 & 0 & -3 \end{pmatrix} & (c) \begin{pmatrix} 3 & -1 & -1 \\ 0 & 2 & -1 \\ 1 & -1 & 3 \end{pmatrix} \\ (d) \begin{pmatrix} 2 & -8 & -4 \\ 0 & -2 & -2 \\ 0 & 8 & 6 \end{pmatrix} & (e) \begin{pmatrix} 4 & -1 & -1 \\ 2 & -1 & -2 \\ -3 & 7 & 6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵⁴Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵⁵

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -2 & 2 & -2 \\ -5 & 6 & -7 \\ -1 & 2 & -3 \end{pmatrix} & (b) \begin{pmatrix} 1 & -6 & -3 \\ 0 & -1 & -1 \\ 0 & 2 & 2 \end{pmatrix} & (c) \begin{pmatrix} 4 & -1 & -1 \\ -1 & 4 & 1 \\ 4 & -5 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} 2 & 1 & 2 \\ 1 & 2 & -2 \\ -1 & 1 & 5 \end{pmatrix} & (e) \begin{pmatrix} 2 & 2 & 1 \\ 3 & 0 & -3 \\ -4 & 5 & 7 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵⁵Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵⁶

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -2 & 3 & 3 \\ 2 & -1 & 1 \\ -4 & 4 & 2 \end{pmatrix} & (b) \begin{pmatrix} 2 & 2 & 0 \\ 0 & 1 & 0 \\ -2 & -4 & 1 \end{pmatrix} & (c) \begin{pmatrix} 2 & 0 & 3 \\ 3 & -1 & 3 \\ 6 & -6 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} 7 & -4 & -2 \\ 4 & -1 & -2 \\ 0 & 0 & 3 \end{pmatrix} & (e) \begin{pmatrix} -1 & 1 & 2 \\ 2 & -2 & 5 \\ -2 & -1 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵⁶Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵⁷

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -2 & 3 & 3 \\ 4 & -3 & 0 \\ -6 & 6 & 3 \end{pmatrix} & (b) \begin{pmatrix} -2 & 4 & 0 \\ 0 & 2 & 0 \\ -4 & 4 & 2 \end{pmatrix} & (c) \begin{pmatrix} -2 & -6 & 3 \\ 4 & 0 & 5 \\ 2 & 4 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 2 & 2 \\ -1 & -4 & -1 \\ -1 & -1 & -4 \end{pmatrix} & (e) \begin{pmatrix} 0 & 2 & 4 \\ -1 & 0 & 1 \\ -2 & -2 & -6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵⁷Date: Dec 19, 2009; Dead Line : 27 Dec 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵⁸

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -1 & -3 & -3 \\ 1 & 3 & 3 \\ 1 & 1 & -1 \end{pmatrix} & (b) \begin{pmatrix} 2 & 0 & 0 \\ 8 & -2 & -4 \\ -4 & 2 & 4 \end{pmatrix} & (c) \begin{pmatrix} -4 & -8 & -7 \\ 1 & -3 & 1 \\ 3 & 7 & 6 \end{pmatrix} \\ (d) \begin{pmatrix} 1 & -3 & 6 \\ -1 & -1 & -2 \\ -2 & 2 & -6 \end{pmatrix} & (e) \begin{pmatrix} -4 & -2 & 6 \\ -4 & -6 & 6 \\ -3 & -3 & 4 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵⁸Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁵⁹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -1 & -3 & 1 \\ 1 & -1 & 1 \\ 1 & 3 & -1 \end{pmatrix} & (b) \begin{pmatrix} 1 & -1 & 0 \\ 0 & 2 & 0 \\ -1 & -1 & 2 \end{pmatrix} & (c) \begin{pmatrix} -3 & -2 & 3 \\ -2 & -3 & 3 \\ -4 & -8 & 7 \end{pmatrix} \\ (d) \begin{pmatrix} -4 & 2 & -2 \\ 0 & -2 & 0 \\ 2 & -2 & 0 \end{pmatrix} & (e) \begin{pmatrix} -6 & 4 & 4 \\ -2 & 0 & 2 \\ -1 & 0 & 0 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁵⁹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁶⁰

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -1 & -2 & -4 \\ 3 & 4 & 4 \\ 3 & 3 & 5 \end{pmatrix} & (b) \begin{pmatrix} 6 & -4 & 1 \\ 3 & -1 & 1 \\ 0 & 0 & 3 \end{pmatrix} & (c) \begin{pmatrix} -1 & 0 & 2 \\ 0 & -3 & 6 \\ 0 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 1 & 0 \\ -1 & -3 & 0 \\ 1 & 1 & -2 \end{pmatrix} & (e) \begin{pmatrix} 4 & 0 & -6 \\ 3 & -2 & -3 \\ 8 & -4 & -8 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁶⁰Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁶¹

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -1 & -2 & -2 \\ -1 & -2 & -2 \\ 1 & -1 & -1 \end{pmatrix} & (b) \begin{pmatrix} 3 & -2 & -2 \\ 0 & 3 & 0 \\ 0 & -2 & 1 \end{pmatrix} & (c) \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 1 \\ -1 & -1 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & 2 & -2 \\ 0 & -2 & 1 \\ 0 & -1 & 0 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & -3 \\ -1 & 0 & 1 \\ 4 & -2 & -5 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁶¹Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁶²

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -1 & -1 & -3 \\ -3 & -3 & 3 \\ -7 & -7 & 3 \end{pmatrix} & (b) \begin{pmatrix} 5 & 0 & 2 \\ 2 & 3 & 2 \\ -4 & 0 & -1 \end{pmatrix} & (c) \begin{pmatrix} 3 & -1 & -1 \\ 0 & 2 & -1 \\ 1 & -1 & 3 \end{pmatrix} \\ (d) \begin{pmatrix} -3 & 4 & 0 \\ -1 & 1 & 0 \\ -2 & 4 & -1 \end{pmatrix} & (e) \begin{pmatrix} 0 & 1 & 0 \\ -1 & 2 & 0 \\ 2 & -3 & 1 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁶²Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁶³

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -1 & -1 & -3 \\ 2 & 2 & 3 \\ 2 & 2 & 3 \end{pmatrix} & (b) \begin{pmatrix} 1 & -1 & 0 \\ 2 & 4 & 0 \\ 4 & 2 & 3 \end{pmatrix} & (c) \begin{pmatrix} 4 & -1 & -1 \\ -1 & 4 & 1 \\ 4 & -5 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} -3 & 4 & 0 \\ -1 & 1 & 0 \\ -3 & 6 & -1 \end{pmatrix} & (e) \begin{pmatrix} 2 & -1 & 0 \\ -2 & 1 & -1 \\ 2 & 3 & 3 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁶³Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁶⁴

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -1 & 0 & -8 \\ 4 & 3 & 8 \\ -4 & -4 & 7 \end{pmatrix} & (b) \begin{pmatrix} -3 & 4 & 4 \\ 0 & -3 & 0 \\ 0 & 2 & -1 \end{pmatrix} & (c) \begin{pmatrix} 2 & 0 & 3 \\ 3 & -1 & 3 \\ 6 & -6 & 2 \end{pmatrix} \\ (d) \begin{pmatrix} -1 & -8 & 4 \\ 0 & -5 & 2 \\ 0 & -8 & 3 \end{pmatrix} & (e) \begin{pmatrix} -4 & -6 & -3 \\ 8 & 8 & 4 \\ 0 & 3 & 2 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁶⁴Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

UNIVERSITY OF HYDERABAD
School of Physics

M.Sc.Physics II Semester

Dec 2009-Apr 2010

Quantum Mechanics-I

Set-I :: Home Work⁶⁵

EIGENVALUES AND EIGENVECTORS

[1] Find eigenvalues and normalized eigenvectors of the following matrices and verify the following statements.

- The first matrix has three distinct eigenvalues.
- The second matrix has two distinct characteristic roots and three linearly independent eigenvectors.
- the third matrix has two different eigenvalues and only two linearly independent eigenvectors.
- All the three eigenvalues of the fourth matrix are equal and it has only two linearly independent eigenvectors.
- All the three eigenvalues of the fifth matrix are equal and it has only one linearly independent eigenvector.

$$\begin{array}{lll} (a) \begin{pmatrix} -1 & 0 & -6 \\ 3 & 2 & 6 \\ -3 & -3 & 5 \end{pmatrix} & (b) \begin{pmatrix} 0 & 0 & 3 \\ -2 & -2 & -3 \\ -2 & 0 & -5 \end{pmatrix} & (c) \begin{pmatrix} -2 & -6 & 3 \\ 4 & 0 & 5 \\ 2 & 4 & -1 \end{pmatrix} \\ (d) \begin{pmatrix} 3 & -4 & 0 \\ 1 & -1 & 0 \\ 2 & -4 & 1 \end{pmatrix} & (e) \begin{pmatrix} 4 & -1 & -1 \\ 2 & -1 & -2 \\ -3 & 7 & 6 \end{pmatrix} & \end{array}$$

[2] Give an example of a 3×3 matrix which has three linearly independent eigenvectors and all three eigenvectors correspond to eigenvalue 1.

[3] Which of the above five matrices can be diagonalized? Give reasons to support your answer.

⁶⁵Date: Dec 19, 2009; Dead Line : 27 Dec 27 2009

ANSWERS (01):

(a) The eigenvalues are $\{-2, 1, 0\}$ and the eigenvector(s) are

$$\{-2, \{3, -2, 1\}\} \quad \{1, \{1, -1, 1\}\} \quad \{0, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{-3, -3, -1\}$ and the eigenvector(s) are

$$\{-3, \{0, -1, 1\}\} \quad \{-3, \{1, 0, 0\}\} \quad \{-1, \{2, 0, 1\}\}$$

(c) The eigenvalues are $\{-2, -2, 1\}$ and the eigenvector(s) are

$$\{-2, \{-3, 1, 2\}\} \quad \{1, \{-1, 1, 1\}\}$$

(d) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 0, 1\}\} \quad \{-3, \{-1, 1, 0\}\}$$

(e) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 2, 0\}\}$$

ANSWERS (02):

(a) The eigenvalues are $\{6, 1, 0\}$ and the eigenvector(s) are

$$\{6, \{1, -1, 1\}\} \quad \{1, \{-1, 2, 1\}\} \quad \{0, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{-3, -2, -2\}$ and the eigenvector(s) are

$$\{-3, \{-1, 1, 1\}\} \quad \{-2, \{-3, 0, 2\}\} \quad \{-2, \{0, 1, 0\}\}$$

(c) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{-2, \{-3, -1, 2\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-2, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, -1, 1\}\}$$

ANSWERS (03):

(a) The eigenvalues are $\{3, 2, 1\}$ and the eigenvector(s) are

$$\{3, \{-3, 2, 2\}\} \quad \{2, \{-1, 0, 1\}\} \quad \{1, \{-3, 1, 2\}\}$$

(b) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{-2, \{-1, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{-1, 1, 1\}$ and the eigenvector(s) are

$$\{-1, \{2, 1, 2\}\} \quad \{1, \{1, 1, 2\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 1, 0\}\}$$

ANSWERS (04):

(a) The eigenvalues are $\{-5, -3, 0\}$ and the eigenvector(s) are

$$\{-5, \{-5, -3, 4\}\} \quad \{-3, \{-2, -1, 2\}\} \quad \{0, \{-1, 0, 1\}\}$$

(b) The eigenvalues are $\{-3, -3, -1\}$ and the eigenvector(s) are

$$\{-3, \{1, 0, 1\}\} \quad \{-3, \{0, 1, 0\}\} \quad \{-1, \{2, 1, 1\}\}$$

(c) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{0, 0, 1\}\} \quad \{-2, \{-1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 1\}\}$$

ANSWERS (05):

(a) The eigenvalues are $\{4, -1, 0\}$ and the eigenvector(s) are

$$\{4, \{1, -1, 1\}\} \quad \{-1, \{-1, 1, 0\}\} \quad \{0, \{-1, 2, 1\}\}$$

(b) The eigenvalues are $\{-3, -1, -1\}$ and the eigenvector(s) are

$$\{-3, \{2, 1, 1\}\} \quad \{-1, \{1, 0, 1\}\} \quad \{-1, \{0, 1, 0\}\}$$

(c) The eigenvalues are $\{2, 1, 1\}$ and the eigenvector(s) are

$$\{2, \{-1, 1, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 1, 1\}\} \quad \{-1, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 2\}\}$$

ANSWERS (06):

(a) The eigenvalues are $\{-4, 1, 0\}$ and the eigenvector(s) are

$$\{-4, \{-1, 1, 0\}\} \quad \{1, \{-1, 1, 1\}\} \quad \{0, \{-2, 3, 1\}\}$$

(b) The eigenvalues are $\{-2, -2, -1\}$ and the eigenvector(s) are

$$\{-2, \{1, 0, 1\}\} \quad \{-2, \{-1, 2, 0\}\} \quad \{-1, \{2, -3, 1\}\}$$

(c) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 1\}\}$$

ANSWERS (07):

(a) The eigenvalues are $\{7, 2, 1\}$ and the eigenvector(s) are

$$\{7, \{1, -1, 1\}\} \quad \{2, \{-1, 1, 0\}\} \quad \{1, \{3, -2, 1\}\}$$

(b) The eigenvalues are $\{-2, -1, -1\}$ and the eigenvector(s) are

$$\{-2, \{3, 1, 0\}\} \quad \{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(c) The eigenvalues are $\{3, 2, 2\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{2, \{1, -1, 3\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\}$$

ANSWERS (08):

(a) The eigenvalues are $\{3, -1, 0\}$ and the eigenvector(s) are

$$\{3, \{1, -1, 1\}\} \quad \{-1, \{3, -2, 1\}\} \quad \{0, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{3, -1, -1\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{-1, \{1, 0, 2\}\} \quad \{-1, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{2, 2, -1\}$ and the eigenvector(s) are

$$\{2, \{1, 1, 0\}\} \quad \{-1, \{-2, -1, 2\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 1, 2\}\} \quad \{-1, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (09):

(a) The eigenvalues are $\{3, -1, 0\}$ and the eigenvector(s) are

$$\{3, \{1, -1, 1\}\} \quad \{-1, \{-1, 1, 0\}\} \quad \{0, \{-1, 2, 1\}\}$$

S

(b) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{-3, 0, 1\}\} \quad \{-1, \{1, 1, 0\}\} \quad \{0, \{-1, 1, 1\}\}$$

(c) The eigenvalues are $\{-2, -2, 1\}$ and the eigenvector(s) are

$$\{-2, \{-3, 1, 2\}\} \quad \{1, \{-1, 1, 1\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\} \quad \{1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (10):

(a) The eigenvalues are $\{-4, -3, 0\}$ and the eigenvector(s) are

$$\{-4, \{-1, 0, 1\}\} \quad \{-3, \{-2, 1, 3\}\} \quad \{0, \{0, 1, 0\}\}$$

(b) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{0, 3, 1\}\} \quad \{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(c) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{-2, \{-3, -1, 2\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{2, 0, 1\}\} \quad \{1, \{0, 1, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$

ANSWERS (11):

(a) The eigenvalues are $\{2, -1, 0\}$ and the eigenvector(s) are

$$\{2, \{-2, -2, 1\}\} \quad \{-1, \{1, 1, 0\}\} \quad \{0, \{-7, -6, 2\}\}$$

(b) The eigenvalues are $\{1, 1, 0\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\} \quad \{1, \{1, 1, 0\}\} \quad \{0, \{2, 1, 0\}\}$$

(c) The eigenvalues are $\{-1, 1, 1\}$ and the eigenvector(s) are

$$\{-1, \{2, 1, 2\}\} \quad \{1, \{1, 1, 2\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\} \quad \{1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$

ANSWERS (12):

(a) The eigenvalues are $\{5, 1, 0\}$ and the eigenvector(s) are

$$\{5, \{-2, -2, 1\}\} \quad \{1, \{-2, -1, 1\}\} \quad \{0, \{-1, 0, 1\}\}$$

(b) The eigenvalues are $\{-3, -3, 1\}$ and the eigenvector(s) are

$$\{-3, \{0, 1, 1\}\} \quad \{-3, \{1, 0, 0\}\} \quad \{1, \{-2, 1, 0\}\}$$

(c) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(d) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{0, -1, 2\}\} \quad \{2, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 2, 0\}\}$$

ANSWERS (13):

(a) The eigenvalues are $\{4, -3, 2\}$ and the eigenvector(s) are

$$\{4, \{-1, 1, 0\}\} \quad \{-3, \{1, 0, 1\}\} \quad \{2, \{-1, 1, 1\}\}$$

(b) The eigenvalues are $\{1, 1, 0\}$ and the eigenvector(s) are

$$\{1, \{0, -1, 2\}\} \quad \{1, \{1, 0, 0\}\} \quad \{0, \{-3, -1, 1\}\}$$

(c) The eigenvalues are $\{2, 1, 1\}$ and the eigenvector(s) are

$$\{2, \{-1, 1, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(d) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{2, 0, 1\}\} \quad \{3, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, -1, 1\}\}$$

ANSWERS (14):

(a) The eigenvalues are $\{3, 2, 0\}$ and the eigenvector(s) are

$$\{3, \{1, -1, 1\}\} \quad \{2, \{-1, 1, 0\}\} \quad \{0, \{3, -2, 1\}\}$$

(b) The eigenvalues are $\{2, 1, 1\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\} \quad \{1, \{0, 0, 1\}\} \quad \{1, \{-2, 1, 0\}\}$$

(c) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(d) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 2\}\} \quad \{3, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 1, 0\}\}$$

ANSWERS (15):

(a) The eigenvalues are $\{-5, -3, 1\}$ and the eigenvector(s) are

$$\{-5, \{-1, 0, 1\}\} \quad \{-3, \{-3, 1, 3\}\} \quad \{1, \{-1, 1, 2\}\}$$

(b) The eigenvalues are $\{-2, 2, 2\}$ and the eigenvector(s) are

$$\{-2, \{1, 0, 1\}\} \quad \{2, \{0, 0, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{3, 2, 2\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{2, \{1, -1, 3\}\}$$

(d) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 0, 1\}\} \quad \{-3, \{-1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 1\}\}$$

ANSWERS (16):

(a) The eigenvalues are $\{-2, -1, 0\}$ and the eigenvector(s) are

$$\{-2, \{-1, 1, 0\}\} \quad \{-1, \{3, -2, 2\}\} \quad \{0, \{3, -2, 3\}\}$$

(b) The eigenvalues are $\{2, 2, 0\}$ and the eigenvector(s) are

$$\{2, \{1, 0, 2\}\} \quad \{2, \{1, 2, 0\}\} \quad \{0, \{0, -2, 1\}\}$$

(c) The eigenvalues are $\{2, 2, -1\}$ and the eigenvector(s) are

$$\{2, \{1, 1, 0\}\} \quad \{-1, \{-2, -1, 2\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-2, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 2\}\}$$

ANSWERS (17):

(a) The eigenvalues are $\{-3, -2, 1\}$ and the eigenvector(s) are

$$\{-3, \{-1, 0, 1\}\} \quad \{-2, \{-2, 1, 3\}\} \quad \{1, \{0, 1, 0\}\}$$

(b) The eigenvalues are $\{2, 2, 1\}$ and the eigenvector(s) are

$$\{2, \{0, 0, 1\}\} \quad \{2, \{-1, 1, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(c) The eigenvalues are $\{-2, -2, 1\}$ and the eigenvector(s) are

$$\{-2, \{-3, 1, 2\}\} \quad \{1, \{-1, 1, 1\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 1\}\}$$

ANSWERS (18):

(a) The eigenvalues are $\{5, 1, 0\}$ and the eigenvector(s) are

$$\{5, \{1, 1, 1\}\} \quad \{1, \{0, -1, 1\}\} \quad \{0, \{1, -1, 2\}\}$$

(b) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 3\}\} \quad \{3, \{4, 3, 0\}\} \quad \{2, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{-2, \{-3, -1, 2\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{0, 0, 1\}\} \quad \{-2, \{-1, 1, 0\}\}$$

(e) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\}$$

ANSWERS (19):

(a) The eigenvalues are $\{3, -1, 0\}$ and the eigenvector(s) are

$$\{3, \{-2, -2, 1\}\} \quad \{-1, \{-1, 0, 1\}\} \quad \{0, \{-2, -1, 1\}\}$$

(b) The eigenvalues are $\{3, 3, 1\}$ and the eigenvector(s) are

$$\{3, \{0, -1, 1\}\} \quad \{3, \{1, 0, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(c) The eigenvalues are $\{-1, 1, 1\}$ and the eigenvector(s) are

$$\{-1, \{2, 1, 2\}\} \quad \{1, \{1, 1, 2\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 1, 1\}\} \quad \{-1, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (20):

(a) The eigenvalues are $\{5, 2, 1\}$ and the eigenvector(s) are

$$\{5, \{1, -1, 1\}\} \quad \{2, \{-1, 1, 0\}\} \quad \{1, \{3, -2, 1\}\}$$

(b) The eigenvalues are $\{3, 3, 1\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{3, \{0, 1, 0\}\} \quad \{1, \{-1, -1, 2\}\}$$

(c) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (21):

(a) The eigenvalues are $\{3, -2, 1\}$ and the eigenvector(s) are

$$\{3, \{-1, 2, 1\}\} \quad \{-2, \{-1, 1, 1\}\} \quad \{1, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{0, 0, 1\}\} \quad \{3, \{-1, 2, 0\}\} \quad \{2, \{-1, 1, 2\}\}$$

(c) The eigenvalues are $\{2, 1, 1\}$ and the eigenvector(s) are

$$\{2, \{-1, 1, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$

ANSWERS (22):

(a) The eigenvalues are $\{3, 2, 0\}$ and the eigenvector(s) are

$$\{3, \{-2, 3, 1\}\} \quad \{2, \{-1, 1, 1\}\} \quad \{0, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{-3, -3, -1\}$ and the eigenvector(s) are

$$\{-3, \{0, -1, 1\}\} \quad \{-3, \{1, 0, 0\}\} \quad \{-1, \{2, 0, 1\}\}$$

(c) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 1, 2\}\} \quad \{-1, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$

ANSWERS (23):

(a) The eigenvalues are $\{-3, -1, 0\}$ and the eigenvector(s) are

$$\{-3, \{1, 0, 1\}\} \quad \{-1, \{5, -3, 3\}\} \quad \{0, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{-3, -2, -2\}$ and the eigenvector(s) are

$$\{-3, \{-1, 1, 1\}\} \quad \{-2, \{-3, 0, 2\}\} \quad \{-2, \{0, 1, 0\}\}$$

(c) The eigenvalues are $\{3, 2, 2\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{2, \{1, -1, 3\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\} \quad \{1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 2, 0\}\}$$

ANSWERS (24):

(a) The eigenvalues are $\{-5, -1, 0\}$ and the eigenvector(s) are

$$\{-5, \{-1, 1, 1\}\} \quad \{-1, \{-1, 2, 1\}\} \quad \{0, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{-2, \{-1, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{2, 2, -1\}$ and the eigenvector(s) are

$$\{2, \{1, 1, 0\}\} \quad \{-1, \{-2, -1, 2\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{2, 0, 1\}\} \quad \{1, \{0, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, -1, 1\}\}$$

ANSWERS (25):

(a) The eigenvalues are $\{-3, 1, 0\}$ and the eigenvector(s) are

$$\{-3, \{-1, 0, 1\}\} \quad \{1, \{-1, 1, 2\}\} \quad \{0, \{-3, 2, 6\}\}$$

(b) The eigenvalues are $\{-3, -3, -1\}$ and the eigenvector(s) are

$$\{-3, \{1, 0, 1\}\} \quad \{-3, \{0, 1, 0\}\} \quad \{-1, \{2, 1, 1\}\}$$

(c) The eigenvalues are $\{-2, -2, 1\}$ and the eigenvector(s) are

$$\{-2, \{-3, 1, 2\}\} \quad \{1, \{-1, 1, 1\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\} \quad \{1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 1, 0\}\}$$

ANSWERS (26):

(a) The eigenvalues are $\{-2, 1, 0\}$ and the eigenvector(s) are

$$\{-2, \{-1, 1, 0\}\} \quad \{1, \{-1, 0, 1\}\} \quad \{0, \{-2, -1, 2\}\}$$

(b) The eigenvalues are $\{-3, -1, -1\}$ and the eigenvector(s) are

$$\{-3, \{2, 1, 1\}\} \quad \{-1, \{1, 0, 1\}\} \quad \{-1, \{0, 1, 0\}\}$$

(c) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{-2, \{-3, -1, 2\}\}$$

(d) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{0, -1, 2\}\} \quad \{2, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 1\}\}$$

ANSWERS (27):

(a) The eigenvalues are $\{-3, -1, 0\}$ and the eigenvector(s) are

$$\{-3, \{-1, 1, 0\}\} \quad \{-1, \{-2, 3, 1\}\} \quad \{0, \{-1, 1, 1\}\}$$

(b) The eigenvalues are $\{-2, -2, -1\}$ and the eigenvector(s) are

$$\{-2, \{1, 0, 1\}\} \quad \{-2, \{-1, 2, 0\}\} \quad \{-1, \{2, -3, 1\}\}$$

(c) The eigenvalues are $\{-1, 1, 1\}$ and the eigenvector(s) are

$$\{-1, \{2, 1, 2\}\} \quad \{1, \{1, 1, 2\}\}$$

(d) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{2, 0, 1\}\} \quad \{3, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 2\}\}$$

ANSWERS (28):

(a) The eigenvalues are $\{-3, -1, 0\}$ and the eigenvector(s) are

$$\{-3, \{1, 1, 1\}\} \quad \{-1, \{1, 1, 2\}\} \quad \{0, \{1, 2, 3\}\}$$

(b) The eigenvalues are $\{-2, -1, -1\}$ and the eigenvector(s) are

$$\{-2, \{3, 1, 0\}\} \quad \{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(c) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(d) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 2\}\} \quad \{3, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 1\}\}$$

ANSWERS (29):

(a) The eigenvalues are $\{-2, -1, 0\}$ and the eigenvector(s) are

$$\{-2, \{-3, 3, 1\}\} \quad \{-1, \{-2, 2, 1\}\} \quad \{0, \{0, 1, 0\}\}$$

(b) The eigenvalues are $\{3, -1, -1\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{-1, \{1, 0, 2\}\} \quad \{-1, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{2, 1, 1\}$ and the eigenvector(s) are

$$\{2, \{-1, 1, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(d) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 0, 1\}\} \quad \{-3, \{-1, 1, 0\}\}$$

(e) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\}$$

ANSWERS (30):

(a) The eigenvalues are $\{-4, -2, 0\}$ and the eigenvector(s) are

$$\{-4, \{-1, -1, 1\}\} \quad \{-2, \{1, 1, 0\}\} \quad \{0, \{0, -1, 1\}\}$$

(b) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{-3, 0, 1\}\} \quad \{-1, \{1, 1, 0\}\} \quad \{0, \{-1, 1, 1\}\}$$

(c) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-2, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (31):

(a) The eigenvalues are $\{-4, 3, 0\}$ and the eigenvector(s) are

$$\{-4, \{-1, -1, 1\}\} \quad \{3, \{0, -1, 1\}\} \quad \{0, \{1, 1, 0\}\}$$

(b) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{0, 3, 1\}\} \quad \{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(c) The eigenvalues are $\{3, 2, 2\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{2, \{1, -1, 3\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (32):

(a) The eigenvalues are $\{-4, 1, 0\}$ and the eigenvector(s) are

$$\{-4, \{-1, -1, 1\}\} \quad \{1, \{0, -1, 1\}\} \quad \{0, \{1, 1, 0\}\}$$

(b) The eigenvalues are $\{1, 1, 0\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\} \quad \{1, \{1, 1, 0\}\} \quad \{0, \{2, 1, 0\}\}$$

(c) The eigenvalues are $\{2, 2, -1\}$ and the eigenvector(s) are

$$\{2, \{1, 1, 0\}\} \quad \{-1, \{-2, -1, 2\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{0, 0, 1\}\} \quad \{-2, \{-1, 1, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$

ANSWERS (33):

(a) The eigenvalues are $\{2, -1, 0\}$ and the eigenvector(s) are

$$\{2, \{1, 1, 0\}\} \quad \{-1, \{0, -1, 1\}\} \quad \{0, \{-3, -4, 2\}\}$$

(b) The eigenvalues are $\{-3, -3, 1\}$ and the eigenvector(s) are

$$\{-3, \{0, 1, 1\}\} \quad \{-3, \{1, 0, 0\}\} \quad \{1, \{-2, 1, 0\}\}$$

(c) The eigenvalues are $\{-2, -2, 1\}$ and the eigenvector(s) are

$$\{-2, \{-3, 1, 2\}\} \quad \{1, \{-1, 1, 1\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 1, 1\}\} \quad \{-1, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$

ANSWERS (34):

(a) The eigenvalues are $\{-3, 2, 0\}$ and the eigenvector(s) are

$$\{-3, \{0, -1, 1\}\} \quad \{2, \{1, 1, 0\}\} \quad \{0, \{-3, -4, 2\}\}$$

(b) The eigenvalues are $\{1, 1, 0\}$ and the eigenvector(s) are

$$\{1, \{0, -1, 2\}\} \quad \{1, \{1, 0, 0\}\} \quad \{0, \{-3, -1, 1\}\}$$

(c) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{-2, \{-3, -1, 2\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 2, 0\}\}$$

ANSWERS (35):

(a) The eigenvalues are $\{4, 3, -2\}$ and the eigenvector(s) are

$$\{4, \{-1, 1, 0\}\} \quad \{3, \{-1, 1, 1\}\} \quad \{-2, \{1, 0, 1\}\}$$

(b) The eigenvalues are $\{2, 1, 1\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\} \quad \{1, \{0, 0, 1\}\} \quad \{1, \{-2, 1, 0\}\}$$

(c) The eigenvalues are $\{-1, 1, 1\}$ and the eigenvector(s) are

$$\{-1, \{2, 1, 2\}\} \quad \{1, \{1, 1, 2\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, -1, 1\}\}$$

ANSWERS (36):

(a) The eigenvalues are $\{3, -2, 1\}$ and the eigenvector(s) are

$$\{3, \{-1, 1, 0\}\} \quad \{-2, \{-2, -1, 2\}\} \quad \{1, \{-1, 0, 1\}\}$$

(b) The eigenvalues are $\{-2, 2, 2\}$ and the eigenvector(s) are

$$\{-2, \{1, 0, 1\}\} \quad \{2, \{0, 0, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 1, 2\}\} \quad \{-1, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 1, 0\}\}$$

ANSWERS (37):

(a) The eigenvalues are $\{7, 2, 1\}$ and the eigenvector(s) are

$$\{7, \{-1, 1, 1\}\} \quad \{2, \{-2, 1, 1\}\} \quad \{1, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{2, 2, 0\}$ and the eigenvector(s) are

$$\{2, \{1, 0, 2\}\} \quad \{2, \{1, 2, 0\}\} \quad \{0, \{0, -2, 1\}\}$$

(c) The eigenvalues are $\{2, 1, 1\}$ and the eigenvector(s) are

$$\{2, \{-1, 1, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\} \quad \{1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 1\}\}$$

ANSWERS (38):

(a) The eigenvalues are $\{2, 1, 0\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 1\}\} \quad \{1, \{-1, 1, 0\}\} \quad \{0, \{-2, -1, 2\}\}$$

(b) The eigenvalues are $\{2, 2, 1\}$ and the eigenvector(s) are

$$\{2, \{0, 0, 1\}\} \quad \{2, \{-1, 1, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(c) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{2, 0, 1\}\} \quad \{1, \{0, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 2\}\}$$

ANSWERS (39):

(a) The eigenvalues are $\{5, 1, 0\}$ and the eigenvector(s) are

$$\{5, \{-1, 1, 1\}\} \quad \{1, \{-2, 1, 1\}\} \quad \{0, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 3\}\} \quad \{3, \{4, 3, 0\}\} \quad \{2, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{3, 2, 2\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{2, \{1, -1, 3\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\} \quad \{1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 1\}\}$$

ANSWERS (40):

(a) The eigenvalues are $\{3, 2, 0\}$ and the eigenvector(s) are

$$\{3, \{-1, 2, 3\}\} \quad \{2, \{-3, 5, 6\}\} \quad \{0, \{-1, 1, 1\}\}$$

(b) The eigenvalues are $\{3, 3, 1\}$ and the eigenvector(s) are

$$\{3, \{0, -1, 1\}\} \quad \{3, \{1, 0, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(c) The eigenvalues are $\{2, 2, -1\}$ and the eigenvector(s) are

$$\{2, \{1, 1, 0\}\} \quad \{-1, \{-2, -1, 2\}\}$$

(d) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{0, -1, 2\}\} \quad \{2, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\}$$

ANSWERS (41):

(a) The eigenvalues are $\{5, -2, 1\}$ and the eigenvector(s) are

$$\{5, \{-1, 1, 1\}\} \quad \{-2, \{0, 1, 0\}\} \quad \{1, \{-2, 2, 1\}\}$$

(b) The eigenvalues are $\{3, 3, 1\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{3, \{0, 1, 0\}\} \quad \{1, \{-1, -1, 2\}\}$$

(c) The eigenvalues are $\{-2, -2, 1\}$ and the eigenvector(s) are

$$\{-2, \{-3, 1, 2\}\} \quad \{1, \{-1, 1, 1\}\}$$

(d) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{2, 0, 1\}\} \quad \{3, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (42):

(a) The eigenvalues are $\{-2, -1, 0\}$ and the eigenvector(s) are

$$\{-2, \{1, -1, 2\}\} \quad \{-1, \{1, 1, 1\}\} \quad \{0, \{0, -1, 1\}\}$$

(b) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{0, 0, 1\}\} \quad \{3, \{-1, 2, 0\}\} \quad \{2, \{-1, 1, 2\}\}$$

(c) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{-2, \{-3, -1, 2\}\}$$

(d) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 2\}\} \quad \{3, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (43):

(a) The eigenvalues are $\{-5, -3, 2\}$ and the eigenvector(s) are

$$\{-5, \{0, -1, 1\}\} \quad \{-3, \{-1, -3, 3\}\} \quad \{2, \{1, 1, 0\}\}$$

(b) The eigenvalues are $\{-3, -3, -1\}$ and the eigenvector(s) are

$$\{-3, \{0, -1, 1\}\} \quad \{-3, \{1, 0, 0\}\} \quad \{-1, \{2, 0, 1\}\}$$

(c) The eigenvalues are $\{-1, 1, 1\}$ and the eigenvector(s) are

$$\{-1, \{2, 1, 2\}\} \quad \{1, \{1, 1, 2\}\}$$

(d) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 0, 1\}\} \quad \{-3, \{-1, 1, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$

ANSWERS (44):

(a) The eigenvalues are $\{4, 2, -1\}$ and the eigenvector(s) are

$$\{4, \{-2, -2, 1\}\} \quad \{2, \{-2, -1, 2\}\} \quad \{-1, \{-1, 0, 1\}\}$$

(b) The eigenvalues are $\{-3, -2, -2\}$ and the eigenvector(s) are

$$\{-3, \{-1, 1, 1\}\} \quad \{-2, \{-3, 0, 2\}\} \quad \{-2, \{0, 1, 0\}\}$$

(c) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-2, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$

ANSWERS (45):

(a) The eigenvalues are $\{3, 2, -1\}$ and the eigenvector(s) are

$$\{3, \{-1, 1, 0\}\} \quad \{2, \{-1, 1, 1\}\} \quad \{-1, \{1, 0, 1\}\}$$

(b) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{-2, \{-1, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{2, 1, 1\}$ and the eigenvector(s) are

$$\{2, \{-1, 1, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 2, 0\}\}$$

ANSWERS (46):

(a) The eigenvalues are $\{3, 1, 0\}$ and the eigenvector(s) are

$$\{3, \{-1, 2, 1\}\} \quad \{1, \{-1, 1, 0\}\} \quad \{0, \{-1, 1, 1\}\}$$

(b) The eigenvalues are $\{-3, -3, -1\}$ and the eigenvector(s) are

$$\{-3, \{1, 0, 1\}\} \quad \{-3, \{0, 1, 0\}\} \quad \{-1, \{2, 1, 1\}\}$$

(c) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{0, 0, 1\}\} \quad \{-2, \{-1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, -1, 1\}\}$$

ANSWERS (47):

(a) The eigenvalues are $\{4, 2, -1\}$ and the eigenvector(s) are

$$\{4, \{-1, 1, 1\}\} \quad \{2, \{-2, 3, 1\}\} \quad \{-1, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{-3, -1, -1\}$ and the eigenvector(s) are

$$\{-3, \{2, 1, 1\}\} \quad \{-1, \{1, 0, 1\}\} \quad \{-1, \{0, 1, 0\}\}$$

(c) The eigenvalues are $\{3, 2, 2\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{2, \{1, -1, 3\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 1, 1\}\} \quad \{-1, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 1, 0\}\}$$

ANSWERS (48):

(a) The eigenvalues are $\{4, -2, 1\}$ and the eigenvector(s) are

$$\{4, \{-1, 1, 1\}\} \quad \{-2, \{0, 1, 0\}\} \quad \{1, \{-2, 2, 1\}\}$$

(b) The eigenvalues are $\{-2, -2, -1\}$ and the eigenvector(s) are

$$\{-2, \{1, 0, 1\}\} \quad \{-2, \{-1, 2, 0\}\} \quad \{-1, \{2, -3, 1\}\}$$

(c) The eigenvalues are $\{2, 2, -1\}$ and the eigenvector(s) are

$$\{2, \{1, 1, 0\}\} \quad \{-1, \{-2, -1, 2\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 1\}\}$$

ANSWERS (49):

(a) The eigenvalues are $\{4, -2, 1\}$ and the eigenvector(s) are

$$\{4, \{-1, 1, 1\}\} \quad \{-2, \{-1, 1, 0\}\} \quad \{1, \{-2, 1, 1\}\}$$

(b) The eigenvalues are $\{-2, -1, -1\}$ and the eigenvector(s) are

$$\{-2, \{3, 1, 0\}\} \quad \{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(c) The eigenvalues are $\{-2, -2, 1\}$ and the eigenvector(s) are

$$\{-2, \{-3, 1, 2\}\} \quad \{1, \{-1, 1, 1\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 2\}\}$$

ANSWERS (50):

(a) The eigenvalues are $\{-3, 2, 0\}$ and the eigenvector(s) are

$$\{-3, \{0, 1, 0\}\} \quad \{2, \{-1, 1, 1\}\} \quad \{0, \{-2, 2, 1\}\}$$

(b) The eigenvalues are $\{3, -1, -1\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{-1, \{1, 0, 2\}\} \quad \{-1, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{-2, \{-3, -1, 2\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 1, 2\}\} \quad \{-1, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 1\}\}$$

ANSWERS (51):

(a) The eigenvalues are $\{4, 1, 0\}$ and the eigenvector(s) are

$$\{4, \{0, 1, 0\}\} \quad \{1, \{1, 1, 1\}\} \quad \{0, \{3, 3, 2\}\}$$

(b) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{-3, 0, 1\}\} \quad \{-1, \{1, 1, 0\}\} \quad \{0, \{-1, 1, 1\}\}$$

(c) The eigenvalues are $\{-1, 1, 1\}$ and the eigenvector(s) are

$$\{-1, \{2, 1, 2\}\} \quad \{1, \{1, 1, 2\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\} \quad \{1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\}$$

ANSWERS (52):

(a) The eigenvalues are $\{-2, 1, 0\}$ and the eigenvector(s) are

$$\{-2, \{0, 1, 0\}\} \quad \{1, \{1, 2, 1\}\} \quad \{0, \{3, 6, 2\}\}$$

(b) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{0, 3, 1\}\} \quad \{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(c) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{2, 0, 1\}\} \quad \{1, \{0, 1, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (52):

(a) The eigenvalues are $\{-2, 1, 0\}$ and the eigenvector(s) are

$$\{-2, \{0, 1, 0\}\} \quad \{1, \{1, 2, 1\}\} \quad \{0, \{3, 6, 2\}\}$$

(b) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{0, 3, 1\}\} \quad \{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(c) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{2, 0, 1\}\} \quad \{1, \{0, 1, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (54):

(a) The eigenvalues are $\{-4, 2, 0\}$ and the eigenvector(s) are

$$\{-4, \{-1, 1, 0\}\} \quad \{2, \{-1, 1, 1\}\} \quad \{0, \{-2, 1, 1\}\}$$

(b) The eigenvalues are $\{-3, -3, 1\}$ and the eigenvector(s) are

$$\{-3, \{0, 1, 1\}\} \quad \{-3, \{1, 0, 0\}\} \quad \{1, \{-2, 1, 0\}\}$$

(c) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(d) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{0, -1, 2\}\} \quad \{2, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$

ANSWERS (55):

(a) The eigenvalues are $\{2, -1, 0\}$ and the eigenvector(s) are

$$\{2, \{1, 3, 1\}\} \quad \{-1, \{0, 1, 1\}\} \quad \{0, \{1, 2, 1\}\}$$

(b) The eigenvalues are $\{1, 1, 0\}$ and the eigenvector(s) are

$$\{1, \{0, -1, 2\}\} \quad \{1, \{1, 0, 0\}\} \quad \{0, \{-3, -1, 1\}\}$$

(c) The eigenvalues are $\{3, 2, 2\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{2, \{1, -1, 3\}\}$$

(d) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{2, 0, 1\}\} \quad \{3, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$

ANSWERS (56):

(a) The eigenvalues are $\{-2, 1, 0\}$ and the eigenvector(s) are

$$\{-2, \{0, -1, 1\}\} \quad \{1, \{1, 1, 0\}\} \quad \{0, \{-3, -4, 2\}\}$$

(b) The eigenvalues are $\{2, 1, 1\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\} \quad \{1, \{0, 0, 1\}\} \quad \{1, \{-2, 1, 0\}\}$$

(c) The eigenvalues are $\{2, 2, -1\}$ and the eigenvector(s) are

$$\{2, \{1, 1, 0\}\} \quad \{-1, \{-2, -1, 2\}\}$$

(d) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 2\}\} \quad \{3, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 2, 0\}\}$$

ANSWERS (57):

(a) The eigenvalues are $\{-3, 1, 0\}$ and the eigenvector(s) are

$$\{-3, \{0, -1, 1\}\} \quad \{1, \{1, 1, 0\}\} \quad \{0, \{-3, -4, 2\}\}$$

(b) The eigenvalues are $\{-2, 2, 2\}$ and the eigenvector(s) are

$$\{-2, \{1, 0, 1\}\} \quad \{2, \{0, 0, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{-2, -2, 1\}$ and the eigenvector(s) are

$$\{-2, \{-3, 1, 2\}\} \quad \{1, \{-1, 1, 1\}\}$$

(d) The eigenvalues are $\{-3, -3, -3\}$ and the eigenvector(s) are

$$\{-3, \{-1, 0, 1\}\} \quad \{-3, \{-1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, -1, 1\}\}$$

ANSWERS (58):

(a) The eigenvalues are $\{2, -1, 0\}$ and the eigenvector(s) are

$$\{2, \{-1, 1, 0\}\} \quad \{-1, \{1, -1, 1\}\} \quad \{0, \{3, -2, 1\}\}$$

(b) The eigenvalues are $\{2, 2, 0\}$ and the eigenvector(s) are

$$\{2, \{1, 0, 2\}\} \quad \{2, \{1, 2, 0\}\} \quad \{0, \{0, -2, 1\}\}$$

(c) The eigenvalues are $\{3, -2, -2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{-2, \{-3, -1, 2\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-2, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 1, 0\}\}$$

ANSWERS (59):

(a) The eigenvalues are $\{-2, -1, 0\}$ and the eigenvector(s) are

$$\{-2, \{-1, 0, 1\}\} \quad \{-1, \{-3, 1, 3\}\} \quad \{0, \{-1, 1, 2\}\}$$

(b) The eigenvalues are $\{2, 2, 1\}$ and the eigenvector(s) are

$$\{2, \{0, 0, 1\}\} \quad \{2, \{-1, 1, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(c) The eigenvalues are $\{-1, 1, 1\}$ and the eigenvector(s) are

$$\{-1, \{2, 1, 2\}\} \quad \{1, \{1, 1, 2\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{-1, 0, 1\}\} \quad \{-2, \{1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 1\}\}$$

ANSWERS (60):

(a) The eigenvalues are $\{5, 2, 1\}$ and the eigenvector(s) are

$$\{5, \{-1, 1, 1\}\} \quad \{2, \{-2, 1, 1\}\} \quad \{1, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 3\}\} \quad \{3, \{4, 3, 0\}\} \quad \{2, \{1, 1, 0\}\}$$

(c) The eigenvalues are $\{-1, -1, 0\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 0\}\} \quad \{0, \{2, 2, 1\}\}$$

(d) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{0, 0, 1\}\} \quad \{-2, \{-1, 1, 0\}\}$$

(e) The eigenvalues are $\{-2, -2, -2\}$ and the eigenvector(s) are

$$\{-2, \{2, 1, 2\}\}$$

ANSWERS (61):

(a) The eigenvalues are $\{-3, -1, 0\}$ and the eigenvector(s) are

$$\{-3, \{1, 1, 0\}\} \quad \{-1, \{-1, -1, 1\}\} \quad \{0, \{0, -1, 1\}\}$$

(b) The eigenvalues are $\{3, 3, 1\}$ and the eigenvector(s) are

$$\{3, \{0, -1, 1\}\} \quad \{3, \{1, 0, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(c) The eigenvalues are $\{2, 1, 1\}$ and the eigenvector(s) are

$$\{2, \{-1, 1, 0\}\} \quad \{1, \{1, 0, 1\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 1, 1\}\} \quad \{-1, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{1, 0, 1\}\}$$

ANSWERS (62):

(a) The eigenvalues are $\{-4, 3, 0\}$ and the eigenvector(s) are

$$\{-4, \{1, 0, 1\}\} \quad \{3, \{-1, 1, 1\}\} \quad \{0, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{3, 3, 1\}$ and the eigenvector(s) are

$$\{3, \{-1, 0, 1\}\} \quad \{3, \{0, 1, 0\}\} \quad \{1, \{-1, -1, 2\}\}$$

(c) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{-1, -1, 1\}\} \quad \{2, \{1, 1, 0\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\}$$

ANSWERS (63):

(a) The eigenvalues are $\{3, 1, 0\}$ and the eigenvector(s) are

$$\{3, \{-1, 1, 1\}\} \quad \{1, \{-2, 1, 1\}\} \quad \{0, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{3, 3, 2\}$ and the eigenvector(s) are

$$\{3, \{0, 0, 1\}\} \quad \{3, \{-1, 2, 0\}\} \quad \{2, \{-1, 1, 2\}\}$$

(c) The eigenvalues are $\{3, 2, 2\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\} \quad \{2, \{1, -1, 3\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 0, 1\}\} \quad \{-1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (64):

(a) The eigenvalues are $\{7, 3, -1\}$ and the eigenvector(s) are

$$\{7, \{-1, 1, 1\}\} \quad \{3, \{-2, 3, 1\}\} \quad \{-1, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{-3, -3, -1\}$ and the eigenvector(s) are

$$\{-3, \{0, -1, 1\}\} \quad \{-3, \{1, 0, 0\}\} \quad \{-1, \{2, 0, 1\}\}$$

(c) The eigenvalues are $\{2, 2, -1\}$ and the eigenvector(s) are

$$\{2, \{1, 1, 0\}\} \quad \{-1, \{-2, -1, 2\}\}$$

(d) The eigenvalues are $\{-1, -1, -1\}$ and the eigenvector(s) are

$$\{-1, \{0, 1, 2\}\} \quad \{-1, \{1, 0, 0\}\}$$

(e) The eigenvalues are $\{2, 2, 2\}$ and the eigenvector(s) are

$$\{2, \{-1, 0, 2\}\}$$

ANSWERS (65):

(a) The eigenvalues are $\{5, 2, -1\}$ and the eigenvector(s) are

$$\{5, \{-1, 1, 1\}\} \quad \{2, \{-2, 3, 1\}\} \quad \{-1, \{-1, 1, 0\}\}$$

(b) The eigenvalues are $\{-3, -2, -2\}$ and the eigenvector(s) are

$$\{-3, \{-1, 1, 1\}\} \quad \{-2, \{-3, 0, 2\}\} \quad \{-2, \{0, 1, 0\}\}$$

(c) The eigenvalues are $\{-2, -2, 1\}$ and the eigenvector(s) are

$$\{-2, \{-3, 1, 2\}\} \quad \{1, \{-1, 1, 1\}\}$$

(d) The eigenvalues are $\{1, 1, 1\}$ and the eigenvector(s) are

$$\{1, \{0, 0, 1\}\} \quad \{1, \{2, 1, 0\}\}$$

(e) The eigenvalues are $\{3, 3, 3\}$ and the eigenvector(s) are

$$\{3, \{1, 0, 1\}\}$$